

ROMIVEX™ DIP

MITICIDE & INSECTICIDE

For Commercial Use Only

General Information

Romivex™ DIP is a contact miticide and insecticide used for dip applications to manage mites and soft-bodied insects, such as two-spotted spider mites, aphids, thrips, mealybugs, and whiteflies, on all plants and crops. It has a 0-hour restricted entry interval (REI) and a 0-day preharvest interval (PHI).

ACTIVE INGREDIENTS

Rosemary oil 50%

OTHER INGREDIENTS

Glycerol monooleate, Sorbitol, Lauric acid, Myristic acid, Palmitic acid, Stearic acid, Oleic acid50%

TOTAL: 100%

FIFRA 25(b) Exempt: This product is a minimum-risk pesticide that has not been registered with the United States Environmental Protection Agency because it qualifies for exemption from registration under Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) Section 25(b).

KEEP OUT OF REACH OF CHILDREN • WARNING (AVISO)

FIRST AID

IF IN EYES	Rinse with water. Get medical attention if irritation develops and persists.
IF ON SKIN OR CLOTHING	Wash off with soap and water. Get medical attention if irritation develops and persists.
IF INHALED	Move to fresh air. Call a poison control center or doctor for further treatment advice if symptoms develop or persist.
IF INGESTED	Rinse mouth with clean water. Get medical attention if symptoms occur.

HOTLINE NUMBER

For emergency call the poison control center at 1-800-222-1222. Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information concerning this product, call the National Pesticides Information Center at 1-800-858-7378, 6:30 a.m. to 4:30 p.m. Pacific time (PT), seven days a week.

PRECAUTIONARY STATEMENTS

If you do not understand the label, find someone to explain it to you in detail (Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle). Causes skin irritation. Wear protective clothing/protective gloves/eye protection/face protection.

DIRECTIONS FOR USE

READ THE ENTIRE LABEL – USE IN ACCORDANCE WITH LABEL INSTRUCTIONS

USE SITES: For use on all agricultural plants and crops. Indoor and outdoor, not limited to greenhouses, nurseries, ornamentals, trees, shrubs, flowers, landscapes, bedding plants, tropical plants, seedlings, plugs, and liners, and unrooted cuttings. Romivex™ DIP is not recommended for cut flowers.

THIS PRODUCT IS A CONCENTRATE AND MUST BE DILUTED BEFORE USING.
USE THE DIP DILUTION IMMEDIATELY. DO NOT STORE THE DILUTED SOLUTION

RECOMMENDED APPLICATION RATES

A fresh treatment solution should be prepared for every 3,000 cuttings.

0.32-0.64 fl.oz./1 gal. of water

3.2-6.4 fl.oz./10 gal. of water

0.25 - 0.5% (v/v)

DIP APPLICATION INSTRUCTIONS - SHAKE WELL BEFORE USING - Clean and disinfect the dipping containers. Make a new dip treatment solution by diluting the recommended amount of Romivex DIP in sufficient clean water at room temperature to submerge the cuttings to be treated fully. The use of an adjuvant is not required. Submerge the entire cuttings in the dip treatment solution, gently agitating for about 15 to 60 seconds to wet the front and back of the leaves and ensure full coverage of the cuttings. NOTE: The submersion time is sufficient when water stops bubbling. Finally, remove the cuttings from the dip submergence solution and allow the excess to drain as needed. Do not stack the draining cuttings. The cuttings are ready to plant immediately.

ADDITIONAL SPRAY APPLICATION INSTRUCTIONS - Suggested dilution rates 32-256 fl. oz./100 gallons of water. A minimum of 100 gallons of finished solution to treat 40,000 sq.ft. (1 acre) is recommended. Use standard spray equipment—treatment Interval: Every 5 to 7 days.

COMPATIBILITY OF MIXTURES - Limited compatibility testing has been conducted for Romivex DIP with other commonly used chemistries (i.e., calcium, hormones, fungicides). As such, mixing or use of Romivex DIP with any other products shall be the exclusive risk and responsibility of the user. Consult specific product labels for additional information. Conduct a mix compatibility jar test when mixing Romivex DIP with other products and test the mixture on a small scale before making large-scale applications.

PLANT SAFETY (PHYTOTOXICITY) - Since plant varieties are numerous and may react differently to Romivex DIP, test the product on a small scale before making large-scale applications. Keep cuttings out of direct sunlight and maintain a cool temperature (60-70°F) during treatment. Do not perform dip treatment on plants under stress or when temperatures exceed 90°F.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water or food by storage and disposal. Store in original container in a cool, well-ventilated, dry place (68-80 °F). Do not freeze. Keep container closed when not in use. **CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. **IF EMPTY:** Offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration. **IF PARTLY FILLED:** Call your local waste agency for disposal instructions.

Warranty: The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the face hereof. All statements made concerning this product apply only when used as directed.

Manufactured by:

AgroPlantae® Inc.

2976 S Peach Ave
Fresno, CA 93725
info@agroplantae.com
Made In USA



Net Contents: **275 Gal (1,039.5 Liters)**

Net Weight: **2,887.5 lbs. (1292.5 Kg)**

Weight per Gal @ 68° F: **10.5 lbs. (4.7 Kg)**

F0426

